

Precautionary Statements:

Hazardous to humans and domestic animals. May irritate eyes, nose, throat and skin. May be harmful if absorbed through skin, inhaled or swallowed. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Keep away from fire or sparks.

Environmental Hazards:

This product is toxic to fish, aquatic invertebrates and oysters. Do not apply directly to water.

Storage and Disposal:

Do not store or transport next to animal feed or food. Store in cool, dry place away from direct sunlight. Completely empty contents into the spray can and then destroy bag and dispose of in a certified area.

DO NOT RE-USE CONTAINER**First Aid:**

If in Eyes: Rinse eyes with clean water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Seek medical attention.

If Inhaled: Move person to fresh air.

If Swallowed: Have person sip a glass of water if they are able to swallow. Do not induce vomiting unless told by the doctor. Do not give anything to an unconscious person. Seek medical attention.

If on Skin: Remove contaminated clothing and rinse immediately with clean water for 15-20 minutes. Seek medical attention.

Note to Physician: Probable mucosal damage may contraindicate use of gastric lavage.

Notice: No warranty of any kind expressed or implied is made concerning the use of this product. User assumes all risks and liabilities resulting from handling use or application whether in accordance with directions or not.

Directions for use:

Use ½ - 1 tablespoon per gallon of water

Crop	Disease	Instruction
Citrus (Non-Bearing Nursery)	Brown Rot, Citrus Canker, Greasy Spot, Melanose, Pink Pitting, Scab	Begin application when disease first threatens. Repeat at 30day intervals or as needed
Cucumber	Angular Leaf Spot, Downy Mildew	Apply weekly when plants begin to vine.
Eggplant	Alternaria Blight, Anthracnose Phomopsis	Begin application prior to development of disease symptoms. Repeat sprays at 7 to 10days intervals or as needed depending on disease severity
Pepper	Bacterial Spots	Begin applications when conditions first favour disease development and repeat at 5 to 10days interval or as needed depending on the disease severity.
Tomato	Anthracnose, Bacterial Speck, Bacterial Spot, Early Blight, Gray Leaf Mold, Late Blight, Septoria Leaf Spot	Begin application when disease first threatens. Repeat at 5 to 10days intervals or as needed depending on disease severity.